

CircLigase™ ssDNA Ligase

Cat. Nos. CL4111K and CL4115K

Connect with Epicentre on our blog (epicentral.blogspot.com),
Facebook (facebook.com/EpicentreBio), and Twitter ([@EpicentreBio](https://twitter.com/EpicentreBio)).

1. Introduction

CirLigase™ ssDNA Ligase[†] is a thermostable ATP-dependent ligase that catalyzes intramolecular ligation (i.e., circularization) of single-stranded DNA (ssDNA) substrates that have both a 5′-monophosphate and a 3′-hydroxyl group. Linear ssDNAs of greater than ~15 bases are circularized by CirLigase ssDNA Ligase. Under standard reaction conditions, virtually no linear concatamers or circular concatamers are produced.

Applications

Production of single-stranded DNA templates for rolling-circle replication or rolling-circle transcription experiments.

2. Specifications

Storage: Store only at –20°C in a freezer without a defrost cycle.

Storage Buffer: CirLigase ssDNA Ligase is supplied in a 50% glycerol solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 1 mM dithiothreitol (DTT), and 0.1% Triton® X-100.

Unit Definition: One unit of CirLigase ssDNA Ligase converts 1 pmol of a linear 5′-monophosphorylated CirLigase Control Oligo (55-mer) into an exonuclease I-resistant circular form in 1 hour at 60°C under standard assay conditions.

CirLigase 10X Reaction Buffer: 0.5 M MOPS (pH 7.5), 0.1 M KCl, 50 mM MgCl₂, and 10 mM DTT.

ATP is added to the reaction to a final concentration of 0.05 mM ATP. For additional optimization, MnCl₂ can be added to a final concentration of 2.5 mM MnCl₂ (see Note 3, Part 4).

Contaminating Activity Assays: CirLigase ssDNA Ligase is free of detectable DNA exonuclease and endonuclease, and RNase activities.

3. Kit Contents

Component	Component Volumes	
	CL4111K (1,000 U)	CL4115K (5,000 U)
CirLigase™ ssDNA Ligase (100 U/μl)	10 μl	50 μl
CirLigase™ 10X Reaction Buffer	50 μl	150 μl
ATP (1 mM)	20 μl	75 μl
MnCl ₂ (50 mM)	20 μl	75 μl
CirLigase™ ssDNA Control Oligo (2 pmol/μl)	10 μl	25 μl
Sterile Water	500 μl	1 ml

4. General Considerations

- Substrate Requirements:** The circularization reaction requires a ssDNA with 5'-phosphate and 3'-hydroxyl groups. The standard CirLigase reaction uses 10 pmol of linear ssDNA.
- Substrate Size:** The ssDNA must be at least ~15 bases in length. Substrates such as single-stranded oligodeoxynucleotides and single-stranded cDNAs can be ligated by the enzyme.
- MnCl₂:** Generally, circularization of ssDNA, such as oligodeoxynucleotides or cDNA, is enhanced by the addition of manganese chloride (MnCl₂) to the reaction to a final reaction concentration of 2.5 mM. A tube of MnCl₂ is included.
- Amount of CirLigase ssDNA Ligase in the Reaction:** The standard reaction conditions (Part 5) use 100 U of the CirLigase enzyme per 20- μ l reaction (~1 μ M enzyme and 0.5 μ M ssDNA substrate). For custom ligation reactions, we recommend maintaining the enzyme concentration in excess of the substrate concentration.
- Sequence Dependence:** Results at Epicentre indicate that the sequence of the ssDNA can strongly influence the efficiency of the circularization reaction.
- Reaction Time:** The CirLigase ssDNA circularization reaction is typically complete in 60 minutes. However, increasing the reaction time may improve the yield of circular DNA with difficult-to-ligate ssDNA substrates.
- Difficult Substrates:** Some ssDNAs are inefficiently circularized in the standard reaction (Part 5). The yield of circular ssDNA from a difficult-to-ligate substrate may be increased by increasing the concentration of CirLigase ssDNA Ligase in the reaction or lengthening the reaction time (see Note 6, above).
- Control Template:** The CirLigase ssDNA Control Oligo provided in the kit is a 55-base oligodeoxynucleotide containing both 5'-phosphate and 3'-hydroxyl ends. Under standard reaction conditions (10 pmol Control Oligo, 100 U CirLigase ssDNA Ligase, 2.5 mM MnCl₂, 1-hour reaction), the linear Control Oligo is converted to circular ssDNA.

5. Kit Procedure

5.A. Ligation Reaction

- Combine the following reaction components:

		Final Concentration
x μ l	Sterile water	---
10 pmol	Single-stranded DNA template	0.5 pmol/ μ l
2 μ l	CirLigase 10X Reaction Buffer	1X
1 μ l	1 mM ATP	50 μ M
1 μ l	50 mM MnCl ₂	2.5 mM
1 μ l	CirLigase ssDNA Ligase (100 U)	5 U/ μ l
20 μ l	Total reaction volume	

- Incubate the reaction at 60°C for 1 hour.

Note: Longer incubation times or larger amounts of CirLigase ssDNA Ligase may improve the yield of circular ssDNA.

3. Incubate the reaction at 80°C for 10 minutes to inactivate the CirLigase ssDNA Ligase.

5.B. Gel Analysis of the Ligation Reaction

The efficiency of a CirLigase ligation reaction can be readily assessed by gel electrophoresis. When ligating oligos, load approximately 1 pmol of linear ssDNA substrate in one gel lane and 2 µl of the standard CirLigase reaction mixture into an adjacent gel lane of a 20% acrylamide/8 M urea denaturing gel. Run the gel and stain with an appropriate DNA-binding dye. The circularized ssDNA product migrates slower (above) the linear ssDNA band (see Fig. 1). In some instances, the adenylated-oligo intermediate can be seen as a band just above the linear ssDNA.

5.C. Removing the Linear ssDNA Template and Adenylated Intermediate from the Reaction

Once the CirLigase reaction has been terminated, the remaining linear ssDNA substrate and linear single-stranded adenylated intermediate can be removed by treatment with Exonuclease I (which digests linear ssDNA) and Exonuclease III (which digests linear double-stranded DNA). The circular ssDNA is resistant to these exonucleases, while the linear ssDNA and adenylated intermediate are digested.

Most linear ssDNA and adenylated intermediate can be eliminated by addition of 20 U of Exonuclease I, followed by incubation at 37°C for 45 minutes.

However, if the linear ssDNA substrate contains hairpins or other secondary structure, treatment with both Exonuclease I and Exonuclease III may be required. We suggest incubating a standard ligation reaction mixture with 10 U of Exonuclease I and 100 U of Exonuclease III at 37°C for 45 minutes.




Figure 1. CirLigase™ ssDNA Ligase converts linear ssDNA into closed circular ssDNA. A 71-nucleotide ssDNA oligo was converted to a circular ssDNA. Lane 1, DNA markers; lane 2, 71-nucleotide linear ssDNA oligo; lane 3, circularization proceeds through an adenylated intermediate; lane 4, closed-circular ssDNA reaction product.

6. Related Products

Cat. #	Concentration	Quantity
CircLigase™ II ssDNA Ligase		
CL9021K		1,000 Units
CL9025K		5,000 Units
Includes: CircLigase™ II ssDNA Ligase, CircLigase™ II 10X Reaction Buffer, 50 mM MnCl ₂ , CircLigase™ ssDNA Control Oligo, Betaine, Sterile Water.		
Exonuclease I, <i>E. coli</i>		
X40501K	20 U/μl	1,000 Units
X40505K	20 U/μl	5,000 Units
X40520K	20 U/μl	20,000 Units
Exonuclease III, <i>E. coli</i>		
EX4405K	200 U/μl	5,000 Units
EX4425K	200 U/μl	25,000 Units
Includes 10X Reaction Buffer.		

CircLigase is a trademark of Epicentre, Madison, Wisconsin.

Triton is a registered trademark of Rohm & Haas, Philadelphia, Pennsylvania.

†CircLigase ssDNA Ligase is covered by issued patents and pending patent applications in the U.S. and certain other countries. The purchase of this product conveys to the buyer the nontransferable right to use the purchased product and components of the product only for not-for-profit research use by the buyer. The buyer cannot sell or otherwise transfer this product or its components to a third party and in particular, no rights are conveyed to the buyer to use the product or its components for commercial use purpose other than for research to gain information that is used by the buyer. Please contact Epicentre related to other uses.

Visit our technical blog: epicentral.blogspot.com