

Poly(A) Polymerase Tailing Kit

Cat. No. PAP5104H

Connect with Epicentre on our blog (epicentral.blogspot.com),
Facebook ([facebook.com/EpicentreBio](https://www.facebook.com/EpicentreBio)), and Twitter ([@EpicentreBio](https://twitter.com/EpicentreBio)).

1. Introduction

The Poly(A) Polymerase Tailing Kit was developed for the rapid and efficient addition of poly(A)-tails to the 3' end of any RNA. Polyadenylation increases the stability of RNA in eukaryotic cells and enhances its ability to be translated after transfection or microinjection.¹⁻³ Poly(A)-tails can also provide priming sites for the synthesis of first-strand cDNA, be used to end-label⁴ or quantitate⁵ mRNA.

The kit features Poly(A) Polymerase which uses ATP as a substrate for template-independent addition of adenosine monophosphates to the 3'-hydroxyl termini of RNA molecules. The standard protocol was designed to produce a poly(A)-tail length of ~150 b on 60 µg of capped RNA. An alternative protocol is also provided for tailing lesser amounts of RNA as well as suggestions on how to adjust the length of the poly(A)-tails generated.

2. Product Specifications

Storage: Store only at -20°C in a freezer without a defrost cycle.

Storage Buffer: Poly(A) Polymerase is supplied in a 50% glycerol solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM dithiothreitol, 0.1 mM EDTA, and 0.1% Triton® X-100.

Unit Definition: One unit of Poly(A) Polymerase converts 1 nmol of ATP into acid-insoluble material in 10 minutes at 37°C under standard assay conditions.

Poly(A) Polymerase 10X Reaction Buffer: 0.5 M Tris-HCl (pH 8.0), 2.5 M NaCl, and 0.1 M MgCl₂. A 10 mM ATP Solution is also provided.

Quality Control: Poly(A) Polymerase is function-tested in a 50-µl reaction containing 50 mM Tris-HCl (pH 8.0), 250 mM NaCl, 10 mM MgCl₂, 5 µg of yeast tRNA, 1 mM ATP, and varying amounts of Poly(A) Polymerase.

Contaminating Activity Assays: All components of the Poly(A) Polymerase Tailing Kit are free of detectable exo- and endonuclease and RNase activities.

3. Kit Contents

Desc.	Concentration	Quantity
Poly(A) Polymerase Tailing Kit Contents		
Poly(A) Polymerase	@ 4 U/µl	100 µl
Poly(A) Polymerase 10X Reaction Buffer		500 µl
10 mM ATP		500 µl
RNase-Free Water		2 ml

4. Related Products

The following products are also available:

- RiboGuard™ RNase Inhibitor

5. Notes on Using the Poly(A) Polymerase Tailing Kit

- Poly(A)-Tail Length:** The standard protocol will generate a poly(A)-tail length of ~150 b. However, the length of poly(A)-tail which can be synthesized by the Poly(A) Polymerase Tailing Kit is dependent upon several reaction parameters. Accordingly, users can customize poly(A)-tails to a desired length by adjusting one or more of these reaction parameters as outlined below.

Assuming all other reaction parameters are kept constant, poly(A)-tail length increases with:

- increasing units of Poly(A) Polymerase (2-16 Units).
- increasing time of incubation (10-60 minutes).
- decreasing amount of substrate RNA (60-1 µg).
- decreasing total reaction volume (100-10 µl).

Customers wishing to customize the length of the poly(A)-tail generated should set-up several test reactions covering a range of the parameter to be changed, in order to find the most appropriate reaction condition for the tail length desired (see Fig. 1).




Figure 1. Customized Poly(A)-tail Lengths. A 1.4-kb transcript was poly(A)-tailed using various reaction conditions to demonstrate the affect on poly(A)-tail lengths. Each lane contains 0.1 µg of the completed poly(A)-tailing reaction product.

Lane 1: RNA MW markers (sizes top to bottom: 4 kb, 3 kb, 2.5 kb, 2 kb, 1.5 kb, 1 kb).

Lane 2: non-poly(A)-tailed RNA.

Lane 3: 60 µg RNA, 16 U enzyme, 37°C for 30 min., 100 µl total rxn volume.

Lane 4: 60 µg RNA, 6 U enzyme, 37°C for 60 min., 100 µl total rxn volume.

Lane 5: 1 µg RNA, 2 U enzyme, 37°C for 60 min., 50 µl total rxn volume.

Lane 6: 1 µg RNA, 2 U enzyme, 37°C for 30 min., 10 µl total rxn volume.

Lane 7: 10 µg RNA, 4 U enzyme, 37°C for 60 min., 50 µl total rxn volume.

Lane 8: 1 µg RNA, 2 U enzyme, 37°C for 60 min., 10 µl total rxn volume.

2. **Reaction Size:** Poly(A) Polymerase tailing reactions can be scaled up or down as desired.
3. **Stopping the Reaction:** A Poly(A) Polymerase tailing reaction may be stopped in a number of different ways, depending on the subsequent uses of the poly(A)-tailed RNA. These include immediate freezing of the completed reaction at -20°C or -70°C , removal of the enzyme via organic solvent extraction (e.g., phenol/chloroform) or chelation of the Mg^{2+} (e.g. EDTA). We do not recommend heat denaturation of the enzyme to stop the reaction due to the potential for RNA thermal degradation.
4. **Addition to an *In vitro* Translation Reaction:** Poly(A)-tailed RNA should be purified (organic extraction/ethanol precipitation, ammonium acetate precipitation, or spin columns) prior to use in *in vitro* translation systems, or *in vivo* experiments.

Standard Protocol

The following protocol is designed to produce a poly(A)-tail length of ~ 150 b on the entire reaction product of a standard *in vitro* transcription capping reaction using Epicentre's AmpliCap-Max High Yield Message Maker Kit. These kits produce up to $60\ \mu\text{g}$ of RNA in a standard reaction with a capping efficiency of 80%. Completed *in vitro* transcription capping reactions may be directly added to the poly(A)-tailing reaction without further purification. See the Notes section if different poly(A)-tail lengths are desired.

1. On ice, combine the following reaction components in the order given:

x	μl	RNase-Free Water
10	μl	Poly(A) Polymerase 10X Reaction Buffer
10	μl	10 mM ATP
2.5	μl	RiboGuard RNase Inhibitor (optional)
20	μl	<i>In vitro</i> Transcription Capping Reaction (60 μg RNA)
2	μl	Poly(A) Polymerase (8 Units)
<hr/>		
100	μl	Total reaction volume

2. Incubate at 37°C for 30 minutes.
(Extending the incubation to 60 minutes results in poly(A)-tails >200 b.)
3. The reaction may be stopped by any one of the following:
 - a) immediate storage at -20°C .
 - b) addition of EDTA to a final concentration of >11 mM.
 - c) phenol/chloroform extraction and salt/alcohol precipitation.

Alternate Protocol

The following protocol is designed to be used as a starting point from which to customize a poly(A)-tailing reaction for use on 1-10 µg of RNA. See the Notes section for the effect of variously altered reaction parameters.

1. On ice, combine the following reaction components in the order given:

x µl	RNase-Free Water
2 µl	Poly(A) Polymerase 10X Reaction Buffer
2 µl	10 mM ATP
0.5 µl	RiboGuard RNase Inhibitor (optional)
1-10 µg	RNA Substrate
1 µl	Poly(A) Polymerase (4 Units)
20 µl	Total reaction volume

2. Incubate at 37°C for 15-20 minutes.
3. The reaction may be stopped by any one of the following:
 - a) immediate storage at -20°C.
 - b) addition of EDTA to a final concentration of >11 mM.
 - c) phenol/chloroform extraction and salt/alcohol precipitation.

8. References

1. Drummond, D.R. *et al.*, (1985) *J. Cell. Biol.* **100**, 1148.
2. Galili, G. *et al.*, (1988) *J. Biol. Chem.* **263**, 5764.
3. Belasco, J. and Brawerman, G. (1993) *Control of Messenger RNA Stability*, Academic Press, San Diego, CA.
4. Lingner, J. and Keller, W. (1993) *Nucleic Acids Res.* **21**, 2917.
5. Krug, M.S. and Berger, S.L. (1987) *Methods Enzymol.* **152**, 262.

AmpliScribe, RiboGuard, and T7-Flash and are trademarks of Epicentre, Madison, Wisconsin.

Triton is a registered trademark of Rohm & Haas, Philadelphia, Pennsylvania.

Visit our technical blog: epicentral.blogspot.com